




## Chemistry-Marking Scheme 2015

Chennai- 56/2/MT

Q.No	Value points	Marks
1	CH <sub>3</sub> -CH <sub>2</sub> -Br.	1
2	1-methoxypropan-2-ol.	1
3	Dispersed phase – Solid , Dispersion medium – Liquid.	1
4	Due to incompletely filled d-orbitals in +2 oxidation state (ie., in Cu <sup>2+</sup> state. )	1
5	3 Faraday / 3F	1
6	As per Raoult's law $p_A = x_A p_A^\circ$ $P_A = p_A^\circ(1 - x_B) = p_A^\circ - p_A^\circ x_B$ $(p_A^\circ - p_A) / p_A^\circ = x_B$ $\Delta p / p_A^\circ = x_B = \frac{W_B M_A}{M_B W_A}$ $M_B = \frac{W_B M_A}{(\Delta p / p_A^\circ) W_A}$	2
7	(i) C <sub>6</sub> H <sub>5</sub> -NH <sub>2</sub> < C <sub>6</sub> H <sub>5</sub> -NH-CH <sub>3</sub> < CH <sub>3</sub> -CH <sub>2</sub> -NH <sub>2</sub> . (ii) CH <sub>3</sub> -NH-CH <sub>3</sub> < CH <sub>3</sub> -CH <sub>2</sub> -NH <sub>2</sub> < C <sub>2</sub> H <sub>5</sub> -OH.	1 1
8	Pentaamminecarbonatocobalt(III) chloride. Ionization isomerism	1 1
8	OR (i) [CuCl <sub>4</sub> ] <sup>2-</sup> (ii) K <sub>2</sub> [Zn(OH) <sub>4</sub> ]	1,1
9	Rate constant is the proportionality constant that relates rate of reaction with concentration of reactants / Rate of the reaction when molar concentration of the reactant becomes unity.  (i) Unit : time <sup>-1</sup> or s <sup>-1</sup> . (ii) Unit : L mol <sup>-1</sup> time <sup>-1</sup> or M <sup>-1</sup> s <sup>-1</sup> .	1  ½ ½
10	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>i)</p> </div> <div style="text-align: center;">  <p>(ii)</p> </div> </div>	1,1
11	$\Delta T_b = K_b m$ $\Delta T_b = K_b (W_B \times 1000 / M_B \times W_A)$ $353.93 - 353.23 = 2.52 \times 1.5 \times 1000 / M_B \times 90$ $M_B = (2.52 \times 1.5 \times 1000) / (0.7 \times 90)$ $= 60.0 \text{ g mol}^{-1}.$	1 1 1

12	(i) Because of $p\pi-p\pi$ multiple bonding in nitrogen (diatomic) which is absent in phosphorus (polymeric / polyatomic).	1												
	(ii) Because of decrease in tendency of $sp^3$ hybridisation from $H_2O$ to $H_2Te$ .	1												
	(iii) Due to their smallest atomic sizes in respective periods, or due to the fact that they have only one electron less than the next noble gas configuration.	1												
13	(i) $CH_3-CH(OH)-CH_3$	1												
	(ii) $CH_3-CH=CH-CH_3$	1												
	(iii) $p-Br-C_6H_4-CO-CH_3$	1												
14	(i) But-1,3-diene, Acrylonitrile; $CH_2=CH-CH=CH_2$ , $CH_2=CH-CN$	$\frac{1}{2} + \frac{1}{2}$												
	(ii) Phenol, Formaldehyde; $C_6H_5OH$ , $HCHO$	$\frac{1}{2} + \frac{1}{2}$												
	(iii) Tetrafluoroethylene; $CF_2=CF_2$ (Note: half mark for name/s and half mark for structure/s)	$\frac{1}{2} + \frac{1}{2}$												
15	(i) Gluconic acid or $COOH-(CHOH)_4-CH_2OH$	1												
	(ii) Peptide linkage or $-NH-CO-$ links	1												
	(iii)	1												
	<table border="1"> <thead> <tr> <th>s.no</th> <th>DNA</th> <th>RNA</th> </tr> </thead> <tbody> <tr> <td>1</td> <td>Sugar is 2-deoxy ribose</td> <td>Sugar is ribose</td> </tr> <tr> <td>2</td> <td>Double helical structure</td> <td>Single stranded structure</td> </tr> <tr> <td></td> <td colspan="2">(or any other one correct difference)</td> </tr> </tbody> </table>		s.no	DNA	RNA	1	Sugar is 2-deoxy ribose	Sugar is ribose	2	Double helical structure	Single stranded structure		(or any other one correct difference)	
	s.no		DNA	RNA										
1	Sugar is 2-deoxy ribose		Sugar is ribose											
2	Double helical structure	Single stranded structure												
	(or any other one correct difference)													
16	(a)(i) $d^2sp^3$ ; Octahedral	$\frac{1}{2} + \frac{1}{2}$												
	(ii) $sp^3$ ; Tetrahedral	$\frac{1}{2} + \frac{1}{2}$												
	(b) 'en', forms chelate.	$\frac{1}{2} + \frac{1}{2}$												
17	(i) Anion vacancies occupied by free electrons in alkali metal halides, (when they have metal excess defects) are called F-centre.	1												
	(ii) When Si or Ge is doped with a trivalent impurity then electron vacancies are created called positive holes which impart electrical conduction. They are called p-type semiconductors.	1												
	(iii) Ferrimagnetism is observed when the magnetic moments are aligned in parallel and antiparallel way in unequal numbers in a substance leading to small net permanent magnetic moment.	1												
18	$\log (k_2 / k_1) = (E_a / 2.303R) (T_2 - T_1) / T_1 T_2$	1												
	$\log [(8 \times 10^{-2}) / (2 \times 10^{-2})] = 20 E_a / 2.303 \times 8.314 \times 300 \times 320$	1												
	$E_a = [\log(4) \times 2.303 \times 8.314 \times 300 \times 320] / 20$	1												
	$E_a = 55336.8 \text{ J mol}^{-1} = 55.34 \text{ kJ mol}^{-1}$ .	1												
19	(i) Due to intramolecular H-bonding in o-nitrophenol /	1												

	<p>p-nitrophenoxide is more stabilized than o-nitrophenoxide due to more delocalization of the negative charge.</p> <p>(ii) The mutual repulsion between bulky alkyl groups is stronger than the I.p-I.p electronic repulsions.</p> <p>(iii) CH<sub>3</sub>ONa is not only nucleophile but also stronger base, thereby leads to elimination reaction of the alkyl halide.</p>	<p>1</p> <p>1</p>
20	<p>(i) <math>C_6H_5NH_2 \xrightarrow{NaNO_2 + HCl / 278K} C_6H_5N_2Cl \xrightarrow{H_3PO_2 + H_2O} C_6H_6</math></p> <p>(ii) <math>CH_3-CONH_2 \xrightarrow{KOH + Br_2} CH_3NH_2</math></p> <p>(iii) <math>C_6H_5NO_2 \xrightarrow{Sn + HCl \text{ or } Fe + HCl} C_6H_5NH_2</math></p> <p style="text-align: center;"><b>OR</b></p> <p>(i) <math>C_2H_5NH_2 + CH_3COCl \xrightarrow{\text{pyridine}} C_2H_5-NHCOCH_3 + HCl</math></p> <p>(ii) <math>C_2H_5NH_2 + C_6H_5SO_2Cl \longrightarrow C_2H_5NH-O_2SC_6H_5 + HCl</math></p> <p>(iii) <math>C_2H_5NH_2 + CHCl_3 + KOH \longrightarrow C_2H_5NC + KCl + H_2O</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
21	<p>(i) In a catalysis process when the reactants and catalyst occur in same phase, the process is called homogeneous catalysis.</p> <p>(ii) The process of settling of colloidal particles forming precipitate is called coagulation.</p> <p>(iii) Polymeric substances or macromolecules when added to suitable solvents form solutions in which the size of the macromolecules may be in colloidal range. Such colloids are known as macromolecular colloids.</p>	<p>1</p> <p>1</p> <p>1</p>
22	<p>(i) The principle of zone refining is that the impurities are more soluble in the melt of metal than in solid state of the metal.</p> <p>(ii) As leaching agent, thereby oxidizing the metal into soluble cyano-complex / [Au(CN)<sub>2</sub>]<sup>-</sup>.</p> <p>(iii) Wrought iron</p>	<p>1</p> <p>1</p> <p>1</p>
23	<p>(i) Social awareness, Health conscious, Caring, empathy, concern. (or any other two values)</p> <p>(ii) (ii) Cartoon display / street display/poster making (or any other correct answer)</p> <p>(iii) Wrong choice and over dose may be harmful.</p> <p>(iv) Saccharin, Aspartame (or any other example)</p>	<p>1</p> <p>1</p> <p>1</p> <p>½ + ½</p>
24	<p>(a)</p> <p>(i) Ce<sup>4+</sup> gets reverted to 3+ oxidation state in aqueous medium hence is a good oxidizing agent / Ce is more stable in +3 oxidation state.</p>	<p>1</p>

24	<p>(ii) Due to very strong metal-metal bonding ( involving large no. of electrons of the d-orbitals)</p> <p>(iii) Mn has maximum no. of unpaired electrons in 3d-orbitals.</p> <p>(b)(i) <math>2\text{MnO}_4^- + 6\text{H}^+ + 5\text{NO}_2^- \longrightarrow 2\text{Mn}^{2+} + 5\text{NO}_3^- + 3\text{H}_2\text{O}</math></p> <p>(ii) <math>\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{Fe}^{2+} \longrightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}</math></p> <p style="text-align: center;"><b>OR</b></p> <p>(a) (i) Due to d-d transitions (involving absorption of energy in visible range ) / unpaired electrons in d- orbitals.</p> <p>(ii) Because Cr is more stable in +3 oxidation state.</p> <p>(iii) Due to stability of <math>5f^0</math>, <math>5f^7</math>, <math>5f^{14}</math> / very small energy difference / comparable energy among 5f, 6d, and 7s orbitals.</p> <p>(b) The overall decrease in atomic and ionic radii from La to Lu (due to poor shielding effect of 4f electrons) is called Lanthanoid contraction. Common oxidation state of Lanthanoids is +3.</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1+1</p>
25	<p>(a) A is <math>\text{C}_6\text{H}_5\text{CHO}</math>; B &amp; C / C &amp; B are <math>\text{C}_6\text{H}_5\text{CH}_2\text{OH}</math> &amp; <math>\text{C}_6\text{H}_5\text{COONa}</math></p> <p>D is <math>\text{C}_6\text{H}_5\text{CH}(\text{OH})\text{CH}_3</math></p> <p>(b) (i) <math>\text{C}_6\text{H}_5\text{-CO-CH}_3</math> forms yellow coloured <math>\text{CHI}_3</math> on heating with <math>\text{I}_2+\text{KOH}</math> but <math>\text{C}_6\text{H}_5\text{-CO-CH}_2\text{-CH}_3</math> does not / equation form.</p> <p>(ii) With neutral <math>\text{FeCl}_3</math>, phenol gives violet coloration but benzoic acid does not. (any other suitable test).</p> <p>(c) </p> <p style="text-align: center;"><b>OR</b></p> <p>(a) (i) <math>\text{CH}_3\text{CH}(\text{OH})\text{CN}</math></p> <p>(ii) <math>\text{CH}_3\text{CH}=\text{N-NH}_2</math></p> <p>(iii) <math>\text{CH}_3\text{CH}_2\text{OH}</math></p> <p>(b) <math>\text{C}_6\text{H}_5\text{-CO-CH}_3 &lt; \text{CH}_3\text{-CO-CH}_3 &lt; \text{CH}_3\text{-CHO}</math></p> <p>(c) <math>\text{CH}_3\text{CHO}</math> gives yellow precipitate of <math>\text{CHI}_3</math> with <math>\text{I}_2 + \text{KOH}</math> but <math>\text{CH}_3\text{CH}_2\text{CHO}</math> does not/ equation form</p>	<p><math>\frac{1}{2} \times 4</math></p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
26	<p><math>E_{\text{Cell}} = (E_{\text{Ag}}^{\circ} - E_{\text{Ni}}^{\circ}) - (0.0591/n) \log[ \text{Ni}^{2+}/(\text{Ag}^+)^2 ]</math></p> <p><math>= (0.80 + 0.25) - 0.02955 \log(10^{-2}/10^{-6})</math></p> <p><math>= 1.05 - 0.0178 = 1.0322 \text{ V}</math></p> <p><math>\Delta G = - n F E_{\text{cell}}</math></p> <p><math>= - 2 \times 96500 \times 1.0322</math></p>	<p>1</p> <p>1</p> <p>1</p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>

	$= -199214 \text{ J mol}^{-1} = -199.2 \text{ kJ mol}^{-1}$	1
	<b>OR</b>	
26	<p>(a) Molar Conductivity (<math>\Lambda_m</math>) = 1000 K / C</p> $= (1000 \times 1.06 \times 10^{-2}) / 0.1$ $= 106 \text{ S cm}^{-2} \text{ mol}^{-1}.$	$\frac{1}{2}$ $\frac{1}{2}$
	<p>Deg. of dissociation (<math>\alpha</math>) = <math>\Lambda_m / \Lambda_m^0</math></p> $= 106 / (50.1 + 76.5)$ $= 0.8373$	1 $\frac{1}{2}$
	<p>(b) Primary battery- non rechargeable whereas secondary battery is chargeable.  Eg: primary battery-dry cell, mercury cell(any one) , secondary battery- lead storage battery, Ni-Cd battery(any one)  (or any other correct example)</p>	$\frac{1}{2}, \frac{1}{2}$ $\frac{1}{2}, \frac{1}{2}$